

Reducing Overdose Risk

You can use any drug safely, including fentanyl and its analogues, as long as you have the right resources!

- **Use slow/less.** Because fentanyl is so strong, a little goes a long way (compared to heroin!) and overdoses can occur quickly, sometimes before you've finished injecting the dose.
- **Try snorting or smoking instead of injecting.** Injecting carries the highest risk for overdose, so shifting to snorting or smoking may help reduce risk. You can still OD by smoking/snorting, especially with fentanyl, so still use slow/less to start
- **Space out doses.** Give yourself time between doses. Fentanyl acts fast, but it's different for everyone, so pay attention to your body!
- **Practice extra caution when using alone.** Have someone check on you if you like using alone. That way, if you do overdose, someone can intervene.
- **In a group? Stagger your use.** Make sure someone is always alert and that at least one person has naloxone/Narcan on them.
- **Test it.** Knowing what's in your drugs can help you decide how much and how best to use them.
- **Always carry Narcan/naloxone.** Be familiar with the signs of an overdose and be prepared to respond with naloxone, no matter what drug you're using.
- **Listen to your body.** Your overall health impacts your overdose risk. Hydrate, eat, and rest as much as possible.

Need naloxone/Narcan, fentanyl test strips, and other harm reduction supplies? *

Syringe Access / Harm Reduction Programming
6th Street Harm Reduction Center
117 6th Street, btw Mission & Minna
Mon, Tues, Thurs, & Fri 9am - 7pm,
Wed 9am - 2pm & 2 - 9pm, Sat 7 - 11pm

SF AIDS Foundation Mobile Sites (look for the white van!)

Mon: Third St & Innes (Bayview) 4 - 6pm
Tues: Bikeway behind Safeway @ Church & Market, near Duboce Ave: 6 - 8pm
Wed: Wiese Alley, off 16th St, btw Mission & Julian: 6 - 8pm
Thurs: Hemlock Alley, off Polk btw Post & Sutter: 7 - 9pm
Fri: 16th & Mission, 7 - 9pm

Glide Harm Reduction Services

330 Ellis, btw Jones & Taylor
Mon & Tues Noon - 5pm
Wed - Fri Noon - 5pm

San Francisco Drug User's Union

149 Turk, btw Jones & Taylor
Sun - Thurs Noon - 7:30pm

Homeless Youth Alliance/SFNE

558 Clayton St, off Haight St
Mon, Wed, & Fri 5:30 - 7:30pm



St. James Infirmary

234 Eddy, btw Jones & Taylor
Thurs & Fri 6 - 8pm



***Syringe access / harm reduction programming is funded to serve our most vulnerable communities, and we need to reserve our resources for them. Please respect the privacy of these program participants if you are not a harm reduction program participant by accessing a free nasal naloxone kit and training here:**

CBHS Pharmacy @ 1380 Howard Street (@ 10th St)
Monday - Friday 8am - 3:30pm

Fentanyl Use & OD Prevention Tips



The DOPE Project is a program of the Harm Reduction Coalition and is funded by the San Francisco Department of Public Health to coordinate San Francisco's response to drug overdose

Please contact us with any questions or to find out where/how to get naloxone in San Francisco

628-225-2801 or marshall@harmreduction.org



Fentanyl Facts

- Fentanyl is a strong synthetic opioid that has been used in clinical settings for decades and is often described as 80 -100 times stronger than morphine, or about 50 times stronger than heroin
- Fentanyl is partly responsible for our country's current overdose crisis, combined with a lack of resources and the criminalization of people who use drugs
- Fentanyl moving through the street market in SF comes in a powder form and can be injected, smoked, or snorted. It has also been found in other drugs in SF, like black tar heroin, meth, cocaine, and "fake" pills
- Fentanyl and fentanyl analogues (some stronger than fentanyl, some weaker) are **NOT** "Narcan resistant." They are opioids and will respond to naloxone/Narcan, should someone be overdosing
- You cannot overdose simply by touching fentanyl. It must be introduced into the blood stream or a mucus membrane in order for someone to feel the effects.

Fentanyl in San Francisco



*Fentanyl is absolutely here!
(And most likely here to stay!)*

It's in other drugs. Currently, most samples of black tar heroin tested throughout the city are coming back positive for the presence of fentanyl. Samples tested in the clinical lab we partner with confirm that fentanyl and some analogues are present in black tar heroin throughout the city. Samples of methamphetamine, cocaine, and counterfeit pills are also testing positive!

It's being sold as-is. Fentanyl is being bought and sold knowingly throughout the city, in the form of a white, gray, or tan powder that can be injected, smoked, or snorted.


Stay vigilant & assume risk!

The street drug supply has always been, and will always be, unpredictable and inconsistent. Assume an overdose risk no matter what drug you're using, and practice as much harm reduction as possible, as consistently as possible: Go slow. Use less. Test your product. If using alone, double down on other strategies. Have someone check on you. Smoke/snort instead of inject. If you're using in a group, stagger your use so someone is always alert. Know the signs of an overdose. Carry naloxone/Narcan and know how to use it. Look out for each other!

Using Fentanyl

Fentanyl is measured in micrograms, and though it can be used safely, the margin of error is much smaller than when using weaker opioids, like heroin.

One microgram = 1 *millionth* of a gram!



1 microgram of fentanyl can be significantly stronger than 1 gram of heroin. Start off using less and pay attention to how it hits you!

Injecting fentanyl carries the highest risk of overdose, so new users may want to start off snorting or smoking to gauge tolerance

Fentanyl is extremely fast-acting, so pay attention. Overdose can occur quickly, and naloxone/Narcan needs to be administered right away followed by rescue breathing if someone stops breathing.

